

Necessary Vaccines for Older Adults


Getting vaccines to protect you from illness is one of the most important things you can do to stay healthy. Vaccines are often injections, sometimes called “shots.”

Vaccines are very safe. Vaccines can have side effects, but for most people, it is more dangerous to risk getting sick. If you have concerns about vaccine side effects or safety, speak to your healthcare provider.

To get your vaccines, you can contact your healthcare provider. You can also ask your local health department for more information on where to get these vaccines. You also may be able to get vaccines at your local neighborhood pharmacy.

There is a government agency called the Centers for Disease Prevention and Control (CDC). The CDC recommends the following vaccines for most older adults.

- **COVID-19 Vaccine**
- **Influenza (Flu) Vaccine**
- **Pneumococcal (Pneumonia) Vaccine**
- **Shingles (Herpes Zoster) Vaccine**
- **Tetanus/Diphtheria Vaccine**



■ COVID-19 Vaccine

What it does: Protects against getting seriously ill from COVID-19.

Who needs it: Everyone 65 years old or older should be up to date with the COVID-19 vaccine.

When to get it: All adults should be up to date on COVID-19 vaccines. You are up to date with your COVID-19 vaccine if you have received the updated Pfizer-BioNTech or Moderna COVID-19 vaccine. The updated vaccines are called “updated” because they protect against both the original virus that causes COVID-19 and the Omicron variant BA.4 and BA.5. The updated vaccines are also called bivalent vaccines.

■ Influenza (Flu) Vaccine

What it does: Protects against serious illness from yearly flu viruses.

Who needs it: All older adults should get a flu vaccine. People with certain conditions should especially get a flu vaccine, since they are at higher risk for serious side effects from the flu. These groups of people include:

- People 65 years old or older
- Nursing home residents
- People with serious health conditions such as heart disease, diabetes, asthma, lung disease, or HIV.
- Caregivers for older adults. This helps them avoid spreading the flu. There are flu vaccines that are specifically for people 65 or older. The CDC recommends that people 65 or older receive any of the flu vaccines specifically for older adults.

You should not get the flu vaccine if you have had an allergic reaction to the flu vaccine in the past. You should also not get a flu vaccine if you have been diagnosed with Guillian-Barre Syndrome within 6 weeks after previously receiving the flu vaccine.

When to get it: New strains of the flu develop all the time. Because of this, you should get the flu vaccine every year. You should get your flu vaccine in the fall.

■ Pneumococcal (Pneumonia) Vaccine

What it does: Protects against serious illness caused by pneumococcal bacteria. Pneumococcal bacteria can cause pneumonia (a serious infection in the lungs) and infections of the blood and brain.

Who needs it: Anyone 65 years or older

When to get it: There are two kinds of pneumococcal vaccines available: Pneumococcal conjugate vaccines (PCV15, or PCV20) and pneumococcal polysaccharide vaccine (PPSV23).

Ask your healthcare provider which vaccine is best for you.

■ Shingles (Herpes Zoster) Vaccine

What it does: Protects you from getting shingles. This vaccine is called Shingrix. It is very effective in reducing the risk of shingles for older adults. It also protects people from developing a side effect that causes chronic pain (called postherpetic neuralgia).

Who needs it: The CDC recommends that people 50 years and older get the Shingrix vaccine. You should get the shingles vaccine even if you have had shingles before or received the older Zostavax shot which is no longer available in the US.

You should not get a Shingrix vaccine if you have had an allergic reaction to any of its ingredients before. You should also not get the vaccine if you currently have shingles. Wait until your symptoms are gone before getting the shingles shot.

When to get it: Shingrix requires two doses. The second dose should be given between 2-6 months after the first dose. If it has been longer than 6 months since the first dose, it's okay to get the second dose at any time.

■ Tetanus/Diphtheria Vaccine

What it does: There are two types of vaccine that protect you from two potentially deadly bacterial infections. One type is called TD. TD protects you from the diseases tetanus and diphtheria. The second type is called Tdap. Tdap protects you from tetanus, diphtheria, and another disease named pertussis (also called “whooping cough”).

Who needs it: Everyone. You should get a one-time dose of the Tdap vaccine if you are 65 or older and have not had the Tdap vaccine previously. This will help protect you and your grandchildren from whooping cough.

When to get it: Once every 10 years.

The CDC also recommends other vaccines for older adults.

These include the measles, mumps, rubella (MMR) vaccine, and vaccines for varicella, hepatitis A and B, and meningococcal disease. You could have a higher risk of getting these diseases if you have certain health problems, occupations, or lifestyles. Ask your healthcare provider if you should get any of these additional vaccines.

For additional information, visit the CDC website at <https://www.cdc.gov/vaccines>.



40 FULTON STREET
SUITE 809
NEW YORK, NY 10038
212.308.1414 TEL
Info@healthinaging.org

The AGS Health in Aging Foundation is dedicated to improving the health, independence, and quality of life of all older people. We aim to empower older adults and caregivers to actively participate in their health care and decision-making.

DISCLAIMER: This information is not intended to diagnose health problems or to take the place of medical advice or care you receive from your physician or other healthcare provider. Always consult your healthcare provider about your medications, symptoms, and health problems. May 2023

©2022 Health in Aging Foundation. All rights reserved.